

(c) *Conditions of use in horses*—(1) *Amount*. Infuse 2 to 2.5 grams per day for 3 to 5 days during estrus.

(2) *Indications for use*. For control of bacterial infections of the uterus (metritis) and as an aid in improving conception in mares with uterine infections caused by bacteria sensitive to gentamicin.

(3) *Limitations*. Do not use in horses intended for human consumption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[71 FR 51727, Aug. 31, 2006, as amended at 78 FR 17597, Mar. 22, 2013; 78 FR 21060, Apr. 9, 2013; 79 FR 10973, Feb. 27, 2014]

§ 529.1044b Gentamicin solution for dipping eggs.

(a) *Specifications*. Each milliliter of solution contains gentamicin sulfate equivalent to 50 milligrams of gentamicin base.

(b) *Sponsors*. See Nos. 000061 and 054925 in § 510.600(c) of this chapter.

(c) *Conditions of use in turkeys*—(1) *Amount*. The drug is added to clean water to provide a dip solution with a gentamicin concentration of 250 to 1,000 parts per million. A concentration of 500 parts per million is recommended. Clean eggs should be held submerged in the gentamicin solution under a vacuum of about 27.5 to 38 centimeters of mercury for 5 minutes followed by additional soaking in gentamicin solution for approximately 10 minutes at atmospheric pressure. Eggs can also be treated by warming them for 3 to 6 hours at approximately 100 °F then immediately submerging them in gentamicin solution maintained at about 40 °F, keeping the eggs submerged for 10 to 15 minutes.

(2) *Indications for use*. As an aid in the reduction or elimination of the following microorganisms from turkey-hatching eggs: *Arizona hinshawii* (paracolon), *Salmonella* Saintpaul, and *Mycoplasma meleagridis*.

(3) *Limitations*. For use in the dipping treatment of turkey-hatching eggs only. Eggs which have been dipped in the drug shall not be used for food.

[40 FR 13881, Mar. 27, 1975, as amended at 52 FR 7833, Mar. 13, 1987; 62 FR 22889, Apr. 28, 1997; 71 FR 13543, Mar. 16, 2006; 79 FR 10973, Feb. 27, 2014]

§ 529.1115 Halothane.

(a) *Specifications*. The drug is a colorless, odorless, nonflammable, nonexplosive, heavy liquid containing 0.01 percent thymol as a preservative.

(b) *Sponsor*. See Nos. 012164 and 054771 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. Two to 5 percent of inhaled atmosphere for induction of anesthesia; 0.5 to 2 percent for maintenance of anesthesia.

(2) *Indications for use*. For nonfood animals for the induction and maintenance of anesthesia.

(3) *Limitations*. Not for use in animals intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 27915, May 22, 1981, as amended at 62 FR 29014, May 29, 1997; 79 FR 10973, Feb. 27, 2014]

§ 529.1150 Hydrogen peroxide.

(a) *Specifications*. Each milliliter of solution contains 396.1 milligrams (mg) hydrogen peroxide (a 35% w/w solution).

(b) *Sponsor*. See No. 050378 in § 510.600(c) of this chapter.

(c) *Conditions of use in finfish*—(1) *Amount*—(i) Freshwater-reared finfish eggs: 500 to 1,000 mg per liter (L) of culture water for 15 minutes in a continuous flow system once per day on consecutive or alternate days until hatch for all coldwater and coolwater species of freshwater-reared finfish eggs or 750 to 1,000 mg/L for 15 minutes in a continuous flow system once per day on consecutive or alternate days until hatch for all warmwater species of freshwater-reared finfish eggs.

(ii) Freshwater-reared salmonids: 100 mg/L for 30 minutes or 50 to 100 mg/L for 60 minutes once per day on alternate days for three treatments in a continuous flow water supply or as a static bath.

(iii) Coolwater species of freshwater-reared finfish fingerlings and adults (except northern pike & paddlefish) and channel catfish fingerlings and adults: 50 to 75 mg/L for 60 minutes once per day on alternate days for three treatments in continuous flow water supply or as a static bath. Coolwater species of freshwater-reared finfish fry (except northern pike, pallid sturgeon &